This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.


Prelim. Amdt dated November 21, 2003 REPLACEMENT SHEETS-DRAWINGS

Delivery of Rho-antagonist tissue adhesive formulation.


1A. Application of tissue adhesive + Rho antagonist to the injured spinal cord with a



Lesion of Corticospinal tract



Corticospinal tract lesion (untreated adult mice)



BEST AVAILABLE COPY

Figure 4

Figure 4a




Figure 4b





Figure 4c



Figure 4d




BEST AVAILABLE COPY

Figure 5: Effect of C3/fibrin treatment on injured corticospinal tract

Regrowing fibres

Regrowing fibres



BEST AVAILABLE COPY

Figure 6: C3/tibrin glue -treated spinal cord

Figure 6A




Figure 6B




Figure 7: Early Functional recovery

Figure 7A







Figure 7b







BBB tests show recovery after C3 treatment

